

Contents lists available at [ScienceDirect](http://ScienceDirect.com)

Cancer Treatment Reviews

journal homepage: www.elsevierhealth.com/journals/ctrv

Controversy

Is there a relation between type of primary melanoma treatment and the development of intralymphatic metastasis? A review of the literature ☆

S. Sloot ^a, M.J. Speijers ^a, E. Bastiaannet ^b, H.J. Hoekstra ^{a,*}^a Department of Surgical Oncology, University Groningen, University Medical Centre Groningen, The Netherlands^b Department of Surgical Oncology, Leiden University Medical Center, Leiden, The Netherlands

ARTICLE INFO

Article history:

Received 29 October 2015

Received in revised form 19 February 2016

Accepted 24 February 2016

Keywords:

Melanoma

Sentinel lymph node biopsy

Lymph node excision

Surgery

Neoplasm metastasis

Recurrence

Review

ABSTRACT

Background: Intralymphatic metastases (ILM) originate from tumor cell emboli entrapped in dermal lymphatics between primary tumor and regional lymph node basin. Because of this origin, sentinel lymph node biopsy (SLNB) might increase ILM by restricting lymph flow.**Methods:** Pubmed, Embase, Cochrane and Medline were searched for articles on ILM between 1980 and September 2014. ILM Incidences were calculated after wide local excision (WLE), excision with elective lymph node dissection (ELND) or therapeutic lymph node dissection (TLND), WLE with SLNB with or without completion lymph node dissection (CLND) and delayed lymph node dissection (DLND) for patients developing nodal metastasis during follow-up.**Results:** In 36 studies, 14,729 patients underwent WLE, 1682 patients WLE/ELND, 362 patients WLE/DLND and 11,201 patients WLE/SLNB. On meta-analysis, ILM occurrence was 3.4% (95% CI 2.8–4.2%). ILM occurred most frequently in the WLE/DLND group (5.5%, 95% CI 3.5–8.7%), followed by WLE/ELND (4.7%, 95% CI 3.1–7.0%), WLE/SLNB (4.5%, 95% CI 3.5–5.7%) and WLE alone (1.9%, 95% CI 1.4–2.7%). 1330 SLNB+ patients were identified and 5783 SLNB– patients. For these groups, on meta-analysis, ILM recurrence was 13.2% (95% CI 10.8–16.2%) and 3.4% (95% CI 2.5–4.5%), respectively ($p = 0.01$).**Conclusion:** In this review SLNB is associated with an increase of ILM with an incidence of 1.9% for WLE vs. 3.4% for SLNB–. Selection bias in this review cannot be excluded. However, ILM occur four times more frequently after SLNB+ than SLNB– procedures and more often after SLNB+/CLND than WLE/DLND or WLE/ELND. ILM should therefore be viewed as a bio-marker of aggressive primary disease.**Synopsis:** Sentinel lymph node biopsy is thought to increase intralymphatic metastasis by restricting lymph flow. This review demonstrates that there is an increase in metastasis, but this result has to be interpreted with caution due to possible selection bias. Aggressive tumor characteristics are likely the cause of this increase.© 2016 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

The behavior of cutaneous melanoma is notoriously unpredictable. 5-year survival rates deteriorate as stage progresses. For stage IA, IB, IIA, IIB, and IIC these survival rates are 97%, 92%, 81%, 70% and 53%, respectively. 5-year survival for locoregional metastasis is 78% (stage IIIA), 59% (stage IIIB) and 40% (stage IIIC) [1]. Once melanoma has metastasized distantly survival is around 15–20%, although these rates are expected to improve upon the

recent introduction of BRAF targeted drugs, checkpoint inhibitors and new generation immunotherapies [2–9]. Long-term follow-up reveals that ulceration and sentinel lymph node status are the strongest predictors for survival [10,11].

The concept of incidence of locoregional metastases increasing with tumor thickness was recognized decades ago [12–14]. Previously, in transit metastases (ITM) and satellite lesions (SL) were considered different entities, but The American Joint Committee on Cancer (AJCC) has classified both ITM and SL in 2002 as intra-lymphatic metastases (ILM) [15]. Historically, SL have been defined to reside within centimeters of the primary tumor location and ITM in the pathway between primary site and regional lymph node basin. The leading hypothesis is that both originate from tumor cell emboli entrapped in dermal lymphatic vessels between primary tumor and regional lymph node basin [16,17]. The appearance of

☆ Presented at the SMR Melanoma Congress, November 17th–20th 2013, Philadelphia, USA.

* Corresponding author at: Department of Surgical Oncology, University Medical Centre Groningen, University of Groningen, Hanzeplein 1, PO Box 30.001, 9700 RB Groningen, The Netherlands. Tel.: +31 50 361 2317.

E-mail address: h.j.hoekstra@umcg.nl (H.J. Hoekstra).

ILM automatically upstages a patient's disease into stage IIIB/IIIC, decreasing 5-year survival to 59% and 40%, respectively [1]. Survival rates for patients with SL alone, SL/ITM, or ITM are identical and similar to that of patients with nodal disease [18]. Scar recurrence, 'true local recurrence', differs in pathophysiology, as these develop from residual cells of the initial melanoma, a result of false-negative margins or microsatellites.

Curative treatment for primary melanoma remains surgery (wide local excision, WLE) [2,19]. Four prospective additional elective lymph node dissection (ELND) trials showed no impact on survival [20–24]. ELND has become redundant after the introduction of the sentinel lymph node biopsy (SLNB) in 1992, which preserves its diagnostic advantage with less morbidity [21–23,25–27]. Patients with a positive SLNB undergo a completion node dissection (CLND). The MSLT-I study showed a small but significant disease-free and melanoma-specific survival benefit in patients with intermediate thickness melanoma (1.2–3.5 mm) and nodal disease following early treatment [28]. Most notably, a melanoma-specific survival improvement of 20% was reported for patients with intermediate thickness melanoma undergoing SLNB as opposed to observation, although the MSLT-I did not show improvement in recurrence free, distant metastasis free and melanoma specific survival for the entire population. The MSLT-II study will answer in the near future whether a CLND is indeed indicated after a positive SLNB [29,30]. Other treatment modalities have included therapeutic lymph node dissection (TLND), for metastatic nodal disease at the time of diagnosis, and delayed lymph node dissection (DLND), for patients developing metastatic nodal disease [31].

SLNB in addition to WLE alone has been suspected of causing ILM by inducing lymphatic stasis or entrapment of melanoma cells [32,33]. Pathophysiology on which this hypothesis is built is that the lymph flow from the skin reaches the nodal basin within minutes, with melanoma cells still in lymphatic channels *en route* to the lymph node basin at the time of SLNB or nodal dissection [33,34]. Estourgie et al. published a fourfold risk of ITM recurrence in SLNB positive patients as compared to SLNB negative patients, thereby raising the question whether surgical treatment of the regional lymph node basin can be responsible for ITM, although the same research group refuted this finding in a larger population [35,36]. Although various authors have studied this phenomenon, most notably Morton et al. in the aforementioned MSLT-I trial and van Poll et al. using data of the Melanoma Institute Australia, a definite answer as to whether the incidence of ILM should be attributed to unfavorable primary tumor characteristics alone or is increased by the SLNB procedure by means of a review of all available data has not yet been published [10,16,28,37,38].

The objective of this review was to provide an extensive body of evidence, answering the question whether ILM frequencies increase after performing SLNB.

Methods

Pubmed, Embase, Cochrane Library and Medline were searched for articles using the terms 'melanoma' and 'recurrence' or 'in transit metastasis' or 'ITM' or 'SL' or 'intralymphatic metastasis' or 'local recurrence' or 'satellite' or 'sentinel node' or 'survival' between January 1980 and September 2014. Articles were excluded if they had not been written in English, if they did not distinguish between a local recurrence and ILM, if incidence for ILM as a first recurrence (FR) was not reported, if studies exclusively reported on SLNB– or SLNB+ or if treatment strategy was unclear. Duplicates, case reports, letters to the editors and case series were excluded. Data regarding ILM as FR derived from our institution's SLNB database (UMCG database) were added to the review.

ITM was classified as recurrent melanoma in the pathway between primary melanoma location and the regional nodal basin, with the lesion more than two or five centimeters from this location, depending on the definition used in the article. All other cutaneous and subcutaneous metastases between the re-excision scar and the location of ITM were classified as SL. As consensus is now that ITM and SL are the same entity, all ITM and SL were combined into one value, 'ILM'.

For all included articles the number of patients with ILM as first recurrence (FR) were calculated per treatment group: for WLE alone, for WLE with ELND, WLE and DLND or TLND and WLE with SLNB. The last group was stratified into tumor-negative SLNB (SLNB–) patients and tumor-positive SLNB (SLNB+) patients undergoing CLND. When assessing risk of ILM as FR, WLE was compared to the WLE/SLNB– group. WLE/SLNB+ was compared to WLE/DLND, WLE/ELND and WLE/TLND groups. As only SLNB+ patients undergo additional CLND, this division groups together the most similar procedures regarding interruption of lymph flow. Additional study characteristics were collected: study design, number of patients, mean/median Breslow thickness, age at diagnosis, and melanoma ulceration status.

Statistical analysis

For a comprehensive review of the data, all data were summarized in tables and analyzed using version 18 SPSS, (IBM, Chicago, Illinois, USA). Descriptive statistics were used to calculate frequencies of ILM for the different treatment strategies. Chi-square tests were used to check for significant differences.

Subsequently, all studies were assigned a weight based on the amount of included patients and entered into a meta-analysis. Meta-analyses were performed stratified for treatment, SLNB results and anatomical localization of the primary tumor. Proportions of ILM and the corresponding 95% CI were calculated and entered in a datasheet. Meta-analyses were performed with the 'metan' module using STATA/SE version 12.0 (StataCorp, College Station, Texas, USA) with the original data as reported in the studies. Pooled ILM proportions and their 95% CI were calculated using a random effects model.

Results

Study characteristics

19,620 studies were identified and assessed according to the inclusion criteria. 36 studies with a total of 33,622 patients were included for analysis (Table 1), including our ongoing academic medical center database (UMCG database). 6 studies were excluded because they exclusively reported on SLNB– or exclusively on SLNB+ patients ($n = 684$ patients) [11,39–43]. Median follow-up ranged from >12 months–11 years. Fifteen out of 36 studies reported mean Breslow depth and 6 reported exclusively median Breslow depth. One study reported Breslow depth using incremental depths [44]. Melanoma ulceration status was reported in 23 studies; in 15 of those data were only available for part of the population. Twelve studies provided treatment/recurrence data on WLE (14,729 patients), 5 on WLE/ELND (1682 patients), 1 on WLE/DLND (362 patients) and 18 on WLE/SLNB (11,201 patients). For the remaining 5648 patients in 7 studies, treatment was not specified. No study reported outcomes exclusively for TLND.

In 23 of the 36 included studies a clear definition of ITM/SL was not provided. ITM was defined as (sub)cutaneous disease recurrence between locoregional lymph node basin and 2, 3 or 5 centimeters from the original scar in $n = 5$, $n = 1$ and $n = 4$ studies, respectively. The remaining 3 studies defined ILM as recurrence

Table 1
Characteristics of included studies.

| No. | Author | Year | No. patients | Age | Follow-up (median) | Breslow (mm, mean) | Ulceration | No. of ILM | % ILM | No. SLNB patients | SN+ | | SN– | |
|-----|---------------------|------|--------------|---|--------------------|------------------------------------|---------------------------|------------|-------|-------------------|-----|-----|------|-----|
| | | | | | | | | | | | Pts | ILM | Pts | ILM |
| 1 | Bagley [12] | 1981 | 103 | NR | >5 years (mean) | NR | NR | 5 | 4.9 | NR | N/A | N/A | N/A | N/A |
| 2 | Janoff [14] | 1982 | 122 | NR | 6.1 years (mean) | NR | NR | 8 | 6.6 | NR | N/A | N/A | N/A | N/A |
| 3 | Roses [13] | 1983 | 658 | NR | 44.8 months (mean) | NR | NR | 15 | 2.3 | NR | N/A | N/A | N/A | N/A |
| 4 | Veronesi [59] | 1991 | 612 | 0–20: 6 21–40: 217 41–50: 159 51–65: 230 | 90 months (mean) | 1.0 | NR | 4 | 0.65 | NR | N/A | N/A | N/A | N/A |
| 5 | Heenan [45] | 1992 | 482 | NR | 5 years (mean) | NR | NR | 7 | 0.62 | NR | N/A | N/A | N/A | N/A |
| 6 | Gadd [60] | 1992 | 1019 | 56 | NR | NR | NR | 89 | 8.7 | NR | N/A | N/A | N/A | N/A |
| 7 | Fusi [44] | 1993 | 1090 | NR | 84 months | <0.75 6% <2.25 38% >2.25 56% | NR | 20 | 1.8 | NR | N/A | N/A | N/A | N/A |
| 8 | Martini [61] | 1994 | 840 | 53.5 | 48 months | 2.3 | NR | 24 | 2.9 | NR | N/A | N/A | N/A | N/A |
| 9 | Karakousis [62] | 1996 | 742 | 48.9 | 92 months (mean) | 2.0 | Present in 25% NR 17 | 47 | 6.3 | NR | N/A | N/A | N/A | N/A |
| 10 | Johnson [63] | 1999 | 306 | 50.6 | 85 months (mean) | NR | NR | 1 | 0.3 | NR | N/A | N/A | N/A | N/A |
| 11 | Borgstein [16] | 1999 | 258 | NR | 27 months | 1.5 (median) | NR | 15 | 4.3 | 258 | 53 | N/A | 205 | N/A |
| 12 | Cohn-Cedermark [46] | 1999 | 2493 | NR* | 11 years | 1.1–2.7 (median)** | NR* | 49 | 1.97 | NR | N/A | N/A | N/A | N/A |
| 13 | Cohn-Cedermark [64] | 2000 | 989 | 51–52 (median) | 11 years | 1.2 (median) | NR | 9 | 0.9 | NR | N/A | N/A | N/A | N/A |
| 14 | Chao [65] | 2002 | 1183 | 52.0 | 16 months | NR | Present in 30% NR 56 | 14 | 1.2 | NR | N/A | N/A | N/A | N/A |
| 15 | Goydos [66] | 2003 | 175 | NR | NR | NR | NR | 14 | 8.0 | 175 | 102 | 14 | 73 | 0 |
| 16 | Estourgie [35] | 2003 | 250 | NR | 72 months | 2.7 | Present in 32% NR 3 | 32 | 10.8 | 250 | 60 | 14 | 190 | 18 |
| 17 | Borgognoni [67] | 2004 | 375 | 55.3 | 35 months | NR | NR | 7 | 1.9 | 375 | 75 | 1 | 300 | 6 |
| 18 | Macripo [68] | 2004 | 274 | 51 (median) | 2.9 years | 1.9 (median) | Present in 8% | 10 | 3.65 | 274 | 46 | 2 | 228 | 8 |
| 19 | Thomas [69] | 2004 | 900 | 57–58 | 60 months | 3.1 (median) | Present in 33% NR 125 | 17 | 1.9 | NR | N/A | N/A | N/A | N/A |
| 20 | Berk [70] | 2005 | 260 | 55 | 29 months | 2.3 | Present in 25% NR 33 | 3 | 1.15 | 260 | 39 | 1 | 221 | 2 |
| 21 | Duprat [71] | 2005 | 240 | 51 (median) | 27.8 months | 1.6 (median) | Present in 30% | 10 | 4.17 | 240 | 42 | N/A | 198 | N/A |
| 22 | Nathansohn [72] | 2005 | 141 | 53 | 41 months | NR | Present in 26% NR 30 | 9 | 6.4 | NR | N/A | N/A | N/A | N/A |
| 23 | Kang [20] | 2005 | 4412 | NR | NR | NR | Present in 9% NR 45.9% | 77 | 1.7 | 1016 | 110 | 9 | 906 | 28 |
| 24 | Van Poll [47] | 2005 | 2018 | 57 | 44 months (mean) | 2.4 | Present in 26% NR 258 | 54 | 2.7 | 754 | 102 | 7 | 652 | 11 |
| 25 | Pawlik [10] | 2005 | 1395 | 51 | 46.8 months | 1.5 (median) | Present in 21% | 86 | 4.9 | 1395 | 234 | 28† | 1136 | 40 |
| 26 | Van Akkooi [73] | 2006 | 262 | NR | 23.3 months | 2.8 | Present in 28% | 11 | 4.2 | 262 | 77 | 7 | 185 | 4 |
| 27 | Cecchi [74] | 2006 | 111 | 53 (median) | 31.5 months | NR | Present in 32% NR 1 | 4 | 3.6 | 111 | 17 | 3 | 94 | 1 |
| 28 | Kretschmer [75] | 2006 | 328 | 60 (median) | 40 months | 2.7 | Present in 34% NR 16 | 25 | 7.6 | NR | N/A | N/A | N/A | N/A |
| 29 | Dalal [76] | 2007 | 1046 | 56 (median) | 36 months (mean) | 2.5 | Present in 28% NR 142 | 50 | 4.8 | 1046 | 163 | 23 | 883 | 27 |
| 30 | Roulin [77] | 2008 | 327 | 54 | 33 months | 2.2 | Present in 27% | 20 | 6.1 | 327 | 74 | 10 | 253 | 10 |
| 31 | UMCG database | 2013 | 589 | 53 | 64.6 months | 3.0 | Present in 35% NR 10 | 45 | 6.1 | 588 | 177 | 30 | 411 | 15 |
| 32 | v/d Broek [78] | 2013 | 305 | 51 | >12 months | 1.6 | Present in 15% NR 20 | 10 | 3.3 | 305 | 54 | 4 | 251 | 6 |

| | | | | | | | | | | | | | | |
|----|----------------|------|------|----------------------|------------------|-----------|---|-----|-----|------|-----|----|----|------|
| 33 | Ribero [79] | 2013 | 1693 | 55.3–57.8 (median)** | 4.8 years | 1.4–2.2** | Present in 82% Yes 549 No 826 Unkn 329 | 92 | 5.4 | 656 | NR | NR | NR | NR |
| 34 | Spillane [49] | 2014 | 1704 | <50.632 >50 1072 | 69 months | NR | | 127 | 7.5 | NR | NR | NR | NR | NR |
| 35 | Martin [50] | 2014 | 80 | 49.9–50.9** | 19.1–33.9 months | 2.9–3.1** | Present in 53% | 12 | 15 | NR | NR | NR | NR | NR |
| 36 | v/d Ploeg [80] | 2014 | 5840 | 56.1–60.2** | 42 months | 2.4–2.5** | Present in 25% NR 781 | 146 | 2.5 | 2909 | 394 | NR | NR | 2515 |

Age and Breslow depth are given as means unless otherwise reported. NR = not reported, classified as number of patients. ILM = intralymphatic metastases.

* Data incomplete.

** Values separately given for two different patient groups.

† SLNB status was split out for 68/86 ILM patients.

within the pathway of lymphatic drainage, between scar and regional nodal basin, and between tumor and nodes, respectively. Seven out of 36 studies distinguished SL from ITM; out of these, 2 studies defined SL and LR as the same entity [13,16,35,45–48].

ILM data review

ILM occurred most frequently in the WLE/DLND group (20/362 patients, 5.5%), followed by WLE/ELND (75/1682 patients, 4.5%), WLE/SLNB (both SLNB+ and SLNB–) (474/11,201 patients, 4.2%), and WLE alone (285/14,729 patients, 1.9%). For the remaining 5648 patients, the occurrence of ILM was not specified according to treatment method. This group includes Spillane et al. and Martin et al, who did provide the amount of patients undergoing SLNB, but did not differentiate recurrence rates for CLND/DLND/TLND and CLND/TLND, respectively [49,50] (Table 2).

Of the 11,201 patients undergoing SLNB, ILM was split out according to tumor status in 6913 patients. Of the SLNB+ group 153/1330 patients (11.5%) developed an ILM as FR versus 176/5783 patients (3.0%) in the SLNB– group. Differences in distribution between the four treatment modalities and differences between SN– and SN+ were statistically significant. ILM as FR after WLE was significantly lower than after WLE/SLNB, WLE/ELND and WLE/DLND (all $p < 0.001$). ILM was significantly lower after WLE/SLNB– compared to WLE/SLNB+ ($p < 0.001$) (Table 3).

Meta-analysis

After review of the data a meta-analysis was performed, with weight assigned to studies based on the amount of included patients. The overall ILM incidence was 3.4% (95% CI 2.8–4.2%). In the meta-analysis, outcomes were similar to the review data with ILM occurring most frequently in the WLE/DLND group (5.5%, 95% CI 3.5–8.7%), followed by WLE/ELND (4.7%, 95% CI 3.1–7.0%), WLE/SLNB (both SLNB+ and SLNB–) (4.5%, 95% CI 3.5–5.7%) and WLE alone (1.9%, 95% CI 1.4–2.7%) (Table 3 and Fig. 1). Of the 11,201 patients undergoing SLNB, ILM was split out according to tumor status in 6913 patients. For the 6913 patients whose SLNB outcome status was reported, ILM recurrence was higher than for the 11,201 patients, i.e. 5.8% (95% CI 4.1–8.3%). For SLNB+ patients, ILM occurrence was higher (13.2%, 95% CI 10.8–16.22%) than for SLNB– patients (3.4%, 95% CI 2.5–4.5%) (Fig. 2).

The WLE group had significantly less ILM recurrence than the SLNB group ($p = 0.02$), but not than WLE/ELND and WLE/DLND ($p = 0.21$ and $p = 0.49$, respectively). SLNB– patients had less recurrence than SLNB+ patients ($p = 0.01$) (Table 3).

Discussion

Background

In this review, 33,622 melanoma patients from 36 studies were analyzed to establish whether performing SLNB on melanoma patients in addition to WLE alone leads to an increase in ILM. This is an ongoing field of discussion in the literature. In fact, Read et al. recently published one of the largest databases so far, ($n = 11,614$) where 505 patients developed ILM as a recurrence at any time during follow-up [51]. ILM percentages were 4.7% and 21.6% for SLNB– and SLNB+ patients, respectively. Numbers were not specified for the 190 patients who developed ILM as FR, which explains partly why the numbers are higher than in our study.

Critics of SLNB have argued that as of yet there is no agreement on adjuvant therapy for node-positive patients and that only 20% of the patients undergoing SLNB will have a positive node [52]. However, nowadays there are new approaches available with tar-

geted and/or immunotherapies that may lead to new adjuvant strategies [53,54]. The argument that no randomized controlled studies have shown a survival advantage for SLNB in node-positive patients has become partly redundant upon publication of the MSLT-I, which shows a (small, but significant) survival advantage for a selective group of patients, i.e. patients with an intermediate thickness melanoma and positive SLNB. Proponents advocate that SLNB is a procedure with a relatively low morbidity and that the current false-negative rate for SLNB performed in reputable institutes is <6%, declining further as experience progresses [55,56].

Results

Based on the results of our meta-analysis, the overall incidence of ILM as FR was 3.4%. Patients who did not undergo any lymph node dissection had the lowest incidence, with 1.9% of patients having ILM recurrence after WLE and 3.4% after SLNB-. ILM occurrence after WLE/DLND and WLE/ELND was slightly higher (4.7% and 5.5%, respectively), but incidence spiked after SLNB+/CLND at 13.2%. For TLND, insufficient data were available.

Differences in ILM occurrence between WLE and WLE/SLNB groups were statistically significant, leading to the conclusion that

Table 2

Reviews classified by treatment, sorted by Breslow thickness for available studies.

| Author | Year | No. patients | No. of ILM | Percentage ILM | Breslow (mm) |
|------------------------------|------|--------------|------------|----------------|--------------|
| <i>WLE (n = 7308)</i> | | | | | |
| Veronesi [59] | 1991 | 612 | 4 | 0.65 | 1.0 mean |
| Van Poll [47] | 2005 | 1035 | 26 | 2.51 | 1.8 mean |
| Martini [61] | 1994 | 840 | 24 | 2.85 | 2.3 mean |
| v/d Ploeg [80] | 2014 | 2931 | 51 | 1.74 | 2.3 mean |
| UMCG database [48] | 2013 | 1 | 0 | 0.00 | 3.0 mean |
| Cohn-Cedermark [64] | 2000 | 989 | 9 | 0.91 | 1.2 (median) |
| Thomas [69] | 2004 | 900 | 17 | 1.89 | 3.1 (median) |
| <i>WLE + ELND (n = 609)</i> | | | | | |
| Karakousis [62] | 1996 | 380 | 27 | 7.11 | 2.0 mean |
| Van Poll [47] | 2005 | 229 | 10 | 4.37 | 3.2 mean |
| <i>WLE + DLND (n = 362)</i> | | | | | |
| Karakousis [62] | 1996 | 362 | 20 | 5.52 | 2.0 mean |
| <i>WLE + SLNB (n = 8868)</i> | | | | | |
| v/d Broek [78] | 2012 | 305 | 6 | 2.0 | 1.6 mean |
| Van Poll [47] | 2005 | 754 | 18 | 2.39 | 1.9 mean |
| Roulin [77] | 2008 | 327 | 20 | 6.12 | 2.2 mean |
| Berk [70] | 2005 | 260 | 3 | 1.15 | 2.3 mean |
| Dalal [76] | 2007 | 1046 | 50 | 4.78 | 2.5 mean |
| v/d Ploeg [80] | 2014 | 2909 | 95 | 3.27 | 2.5 mean |
| Estourgie [35] | 2003 | 250 | 27 | 10.80 | 2.7 mean |
| Van Akkooi [73] | 2006 | 262 | 11 | 4.20 | 2.8 mean |
| UMCG database | 2013 | 588 | 45 | 7.65 | 3.0 mean |
| Duprat [71] | 2005 | 240 | 10 | 4.17 | 1.6 (median) |
| Pawlik [10] | 2005 | 1395 | 86 | 6.16 | 1.5 (median) |
| Borgstein [16] | 1999 | 258 | 11 | 4.26 | 1.5 (median) |
| Macripo [68] | 2004 | 274 | 10 | 3.65 | 1.9 (median) |

NR = not reported, classified as number of patients. ILM = intralymphatic metastases.

* Separate values given for separate treatment groups.

Table 3

Pooled values and total number of ILM in the treatment groups.

| Treatment ^a | Pooled value from meta-analyses | | |
|------------------------|---------------------------------|-------------|-----------------|
| | Estimate | 95% CI | p-value |
| WLE | 1.92 | 1.39–2.66 | Reference value |
| WLE + ELND | 4.67 | 3.10–7.04 | 0.21 |
| WLE + DLND | 5.52 | 3.50–8.70 | 0.49 |
| WLE + SLNB | 4.46 | 3.51–5.67 | 0.02 |
| SN– | 3.35 | 2.52–4.46 | Reference value |
| SN+ | 13.24 | 10.80–16.22 | 0.01 |
| Treatment ^b | Number of ILM | | |
| | Total | ILM (%) | No ILM (%) |
| WLE | 14,729 | 285 (1.9) | 14,444 (98.1) |
| WLE + ELND | 1682 | 75 (4.5) | 1607 (95.5) |
| WLE + DLND | 362 | 20 (5.5) | 342 (94.5) |
| WLE + SLNB | 11,201 | 474 (4.2) | 10,727 (95.8) |
| SN– | 5783 | 176 (3.0) | 5607 (97.0) |
| SN+ | 1330 | 153 (11.5) | 1177 (88.5) |

^a Pooled estimates from the meta-analyses, according to treatment as shown in Figs. 1 and 2.

^b Total number of ILM in the treatment groups for the initial treatments and stratified for SN– and SN+, review data. P-value for differences in distribution (Chi2).

p-value four groups: <0.001

SN– and SN+: p < 0.001




Fig. 1. Pooled percentage of ILM according to treatment.

a sentinel lymph node biopsy alone is associated with an increase in the risk of ILM (from 1.9% to 3.4%, $p = 0.01$).

To test the stasis hypothesis, the most comparable treatment modalities regarding lymph flow disruption are WLE vs. WLE/SLNB– and WLE/SLNB+/CLND vs. WLE/ELND. As metastasis already has occurred in WLE/DLND groups, this is not a good comparator. As ILM incidence according to meta-analysis doubled between WLE vs. WLE/SLNB– and increased almost threefold from 4.7% to 13.2% between WLE/ELND and WLE/SLNB+/CLND groups, ($p < 0.001$), the increase of ILM is unlikely to be due to the increase in lymph stasis. CLND and ELND are comparable in their amount of

lymph flow disruption. This suggests that an aggressive tumor behavior is the main reason for ILM, a statement that is supported by the spike in incidence after SLNB+, which is the patient group with the most aggressive tumor biology.

Limitations

Inevitable to any review, authors use different definitions and inclusion criteria. The level of heterogeneity is considerable, as illustrated in Table 1, where data on patient and tumor characteristics are shown. The inconsistent and varied application of terms




Fig. 2. Pooled percentage of ILM according to SLNB positive or negative result.

as ITM, SL and LR complicate comparisons among trials. Recently some authors have even abandoned the concept of a true local recurrence, merging ITM, SL and local recurrence into locoregional metastasis, leading to considerable data loss [57]. Also, data on mitosis index, Breslow thickness and ulceration status were inconsistent, thus complicating comparisons, necessitating interpreting the results with caution. In general, patients included in SLNB studies have less favorable primary tumor characteristics than patients who undergo WLE alone [58]. Moreover, before introduction of the SLNB technique, patients with less favorable tumor characteristics were to undergo ELND and would therefore not be included in WLE studies. These limitations may account for the difference between this review and the MSLT-I, a prospective study, in which no increase in ILM or local metastasis was reported between biopsy and observation groups ($7.7 \pm 1.0\%$ and $8.4 \pm 1.3\%$, respectively; $p = 0.38$). As we included WLE patients before introduction of SLNB our WLE population would differ from the MSLT-I population.

The percentage of ILM after DLND in our study is lower than expected. This may be due to the small sample size and also due to bias as we only included ILM as FR after DLND. Since these patients have aggressive disease, they may more often progress to distant metastasis instead of locoregional disease.

Summary

This review showed an increase in ILM of 1.5% after only performing a SLNB procedure (ILM 1.9% for WLE vs. 3.4% for SLNB+).

Taking into account the patient groups traditionally included in WLE studies it is difficult to say whether this increase represents an actual increase in ILM recurrence or a selection bias.

The SLNB procedure is the most important prognostic tool in clinical practice, providing a survival benefit in selected SLNB+ patients undergoing CLND and potentially serving as a marker to identify patients for adjuvant therapy. Sentinel lymph node biopsy has been suspected of causing to increase intralymphatic metastasis by restricting lymph flow. This review demonstrates this increase, but this result has to be interpreted with caution due to possible selection bias. As the stasis hypothesis seems to be incorrect based on the data in this study, aggressive tumor characteristics are likely the cause of this increase. We therefore advocate performing SLNB procedures, but to proceed with caution, adhere to the guidelines and not extend the indication area.

Conflict of interest statement

The authors declare no conflict of interest.

Role of the funding source

S. Sloot, MD received a research grant from the Groningen Melanoma and Sarcoma Foundation.

References

- [1] Balch CM, Gershenwald JE, Soong SJ, et al. Final version of 2009 AJCC melanoma staging and classification. *J Clin Oncol* 2009;27:6199–206.
- [2] Sladden MJ, Balch C, Barzilai DA, et al. Surgical excision margins for primary cutaneous melanoma. *Cochrane Database Syst Rev* 2009. CD004835.
- [3] <<http://seer.cancer.gov/statfacts/html/melan.html>>; [accessed 30.03.2013].
- [4] Chapman PB, Hauschild A, Robert C, et al. Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N Engl J Med* 2011;364:2507–16.
- [5] Hodi FS, O'Day SJ, McDermott DF, et al. Improved survival with ipilimumab in patients with metastatic melanoma. *N Engl J Med* 2010;363:711–23.
- [6] Topalian SL, Sznol M, McDermott DF, et al. Survival, durable tumor remission, and long-term safety in patients with advanced melanoma receiving nivolumab. *J Clin Oncol* 2014;32:1020–30.
- [7] Green J, Ariyan C. Update on immunotherapy in melanoma. *Surg Oncol Clin N Am* 2015;24:337–46.
- [8] Robert C, Karaszewska B, Schachter J, et al. Improved overall survival in melanoma with combined dabrafenib and trametinib. *N Engl J Med* 2015;372:30–9.
- [9] Larkin J, Ascierto PA, Dreno B, et al. Combined vemurafenib and cobimetinib in BRAF-mutated melanoma. *N Engl J Med* 2014;371:1867–76.
- [10] Pawlik TM, Ross MI, Johnson MM, et al. Predictors and natural history of in-transit melanoma after sentinel lymphadenectomy. *Ann Surg Oncol* 2005;12:587–96.
- [11] Rossi CR, De Salvo GL, Bonandini E, et al. Factors predictive of nonsentinel lymph node involvement and clinical outcome in melanoma patients with metastatic sentinel lymph node. *Ann Surg Oncol* 2008;15:1202–10.
- [12] Bagley FH, Cady B, Lee A, Legg MA. Changes in clinical presentation and management of malignant melanoma. *Cancer* 1981;47:2126–34.
- [13] Roses DF, Harris MN, Rigel D, Carrey Z, Friedman R, Kopf AW. Local and in-transit metastases following definitive excision for primary cutaneous malignant melanoma. *Ann Surg* 1983;198:65–9.
- [14] Janoff KA, Moseson D, Nohlgren J, Davenport C, Richards C, Fletcher WS. The treatment of state I melanoma of the extremities with regional hyperthermic isolation perfusion. *Ann Surg* 1982;196:316–23.
- [15] Cancer AJCo. Melanoma of the skin staging. 7th ed, 2009.
- [16] Borgstein PJ, Meijer S, van Diest PJ. Are locoregional cutaneous metastases in melanoma predictable? *Ann Surg Oncol* 1999;6:315–21.
- [17] Oashi K, Furukawa H, Nishihara H, et al. Pathophysiological characteristics of melanoma in-transit metastasis in a lymphedema mouse model. *J Invest Dermatol* 2013;133:537–44.
- [18] Hayes AJ, Clark MA, Harries M, Thomas JM. Management of in-transit metastases from cutaneous malignant melanoma. *Br J Surg* 2004;91:673–82.
- [19] Lens MB, Dawes M, Goodacre T, Bishop JA. Excision margins in the treatment of primary cutaneous melanoma: a systematic review of randomized controlled trials comparing narrow vs wide excision. *Arch Surg* 2002;137:1101–5.
- [20] Kang JC, Wanek LA, Essner R, Faries MB, Foshag LJ, Morton DL. Sentinel lymphadenectomy does not increase the incidence of in-transit metastases in primary melanoma. *J Clin Oncol* 2005;23:4764–70.
- [21] Cascinelli N. Margin of resection in the management of primary melanoma. *Semin Surg Oncol* 1998;14(4):272–5.
- [22] Veronesi U, Adamus J, Bandiera DC, et al. Delayed regional lymph node dissection in stage I melanoma of the skin of the lower extremities. *Cancer* 1982;49:2420–30.
- [23] Balch CM, Soong SJ, Bartolucci AA, et al. Efficacy of an elective regional lymph node dissection of 1 to 4 mm thick melanomas for patients 60 years of age and younger. *Ann Surg* 1996;224:255–63 [discussion 63–6].
- [24] Sim FH, Taylor WF, Pritchard DJ, Soule EH. Lymphadenectomy in the management of stage I malignant melanoma: a prospective randomized study. *Mayo Clin Proc* 1986;61:697–705.
- [25] Sim FH, Taylor WF, Ivins JC, Pritchard DJ, Soule EH. A prospective randomized study of the efficacy of routine elective lymphadenectomy in management of malignant melanoma. Preliminary results. *Cancer* 1978;41:948–56.
- [26] Faries MB, Thompson JF, Cochran A, et al. The impact on morbidity and length of stay of early versus delayed complete lymphadenectomy in melanoma: results of the Multicenter Selective Lymphadenectomy Trial (I). *Ann Surg Oncol* 2010;17:3324–9.
- [27] de Vries M, Vonkeman WG, van Ginkel RJ, Hoekstra HJ. Morbidity after inguinal sentinel lymph node biopsy and completion lymph node dissection in patients with cutaneous melanoma. *Eur J Surg Oncol* 2006;32:785–9.
- [28] Morton DL, Thompson JF, Cochran AJ, et al. Final trial report of sentinel-node biopsy versus nodal observation in melanoma. *N Engl J Med* 2014;370:599–609.
- [29] Morton DL. Overview and update of the phase III Multicenter Selective Lymphadenectomy Trials (MSLT-I and MSLT-II) in melanoma. *Clin Exp Metastasis* 2012;29:699–706.
- [30] Leiter U SR, Mauch C, et al. Survival of SLNB-positive melanoma patients with and without complete lymph node dissection: a multicenter, randomized DECOG trial. *J Clin Oncol* 2015;33 [suppl; abstr LBA9002].
- [31] Fisher SR. Elective, therapeutic, and delayed lymph node dissection for malignant melanoma of the head and neck: analysis of 1444 patients from 1970 to 1998. *Laryngoscope* 2002;112:99–110.
- [32] Cascinelli N, Bufalino R, Marolda R, et al. Regional non-nodal metastases of cutaneous melanoma. *Eur J Surg Oncol* 1986;12:175–80.
- [33] Estourgie SH, Nieweg OE, Kroon BB. High incidence of in-transit metastases after sentinel node biopsy in patients with melanoma. *Br J Surg* 2004;91:1370–1.
- [34] Daryanani D, Komdeur R, Hoekstra HJ. Lymphatic entrapment of tumour cells after sentinel lymph-node biopsy for melanoma. *Lancet Oncol* 2000;1:211.
- [35] Estourgie SH, Nieweg OE, Valdes Olmos RA, Hoefnagel CA, Kroon BB. Review and evaluation of sentinel node procedures in 250 melanoma patients with a median follow-up of 6 years. *Ann Surg Oncol* 2003;10:681–8.
- [36] Veenstra HJ, van der Ploeg IM, Wouters MW, Kroon BB, Nieweg OE. Reevaluation of the locoregional recurrence rate in melanoma patients with a positive sentinel node compared to patients with palpable nodal involvement. *Ann Surg Oncol* 2010;17:521–6.
- [37] Morton DL, Thompson JF, Cochran AJ, et al. Sentinel-node biopsy or nodal observation in melanoma. *N Engl J Med* 2006;355:1307–17.
- [38] Pawlik TM, Ross MI, Thompson JF, Eggermont AM, Gershenwald JE. The risk of in-transit melanoma metastasis depends on tumor biology and not the surgical approach to regional lymph nodes. *J Clin Oncol* 2005;23:4588–90.
- [39] Gadd MA, Cosimi AB, Yu J, et al. Outcome of patients with melanoma and histologically negative sentinel lymph nodes. *Arch Surg* 1999;381–7.
- [40] Gad D, Hoiland-Carlson PF, Bartram P, Clemmensen O, Bischoff-Mikkelsen M. Staging patients with cutaneous malignant melanoma by same-day lymphoscintigraphy and sentinel lymph node biopsy: a single-institutional experience with emphasis on recurrence. *J Surg Oncol* 2006;94:94–100.
- [41] De Giorgi V, Leporatti G, Massi D, et al. Outcome of patients with melanoma and histologically negative sentinel lymph nodes: one institution's experience. *Oncology* 2007;73:401–6.
- [42] Roka F, Mastan P, Binder M, et al. Prediction of non-sentinel node status and outcome in sentinel node-positive melanoma patients. *Eur J Surg Oncol* 2008;34:82–8.
- [43] Meyers MO, Yeh JJ, Frank J, et al. Method of detection of initial recurrence of stage II/III cutaneous melanoma: analysis of the utility of follow-up staging. *Ann Surg Oncol* 2009;16:941–7.
- [44] Fusi S, Ariyan S, Sternlicht A. Data on first recurrence after treatment for malignant melanoma in a large patient population. *Plast Reconstr Surg* 1993;91:94–8.
- [45] Heenan PJ, English DR, Holman CD, Armstrong BK. The effects of surgical treatment on survival and local recurrence of cutaneous malignant melanoma. *Cancer* 1992;69:421–6.
- [46] Cohn-Cedermark G, Mansson-Brahme E, Rutqvist LE, Larsson O, Singnorklao T, Ringborg U. Metastatic patterns, clinical outcome, and malignant phenotype in malignant cutaneous melanoma. *Acta Oncol* 1999;38:549–57.
- [47] van Poll D, Thompson JF, Colman MH, et al. A sentinel node biopsy does not increase the incidence of in-transit metastasis in patients with primary cutaneous melanoma. *Ann Surg Oncol* 2005;12:597–608.
- [48] Speijers MJ, Bastiaannet E, Sloot S, Suurmeijer AJ, Hoekstra HJ. Tumor mitotic rate added to the equation: melanoma prognostic factors changed?: a single-institution database study on the prognostic value of tumor mitotic rate for sentinel lymph node status and survival of cutaneous melanoma patients. *Ann Surg Oncol* 2015.
- [49] Spillane AJ, Pasquali S, Haydu LE, Thompson JF. Patterns of recurrence and survival after lymphadenectomy in melanoma patients: clarifying the effects of timing of surgery and lymph node tumor burden. *Ann Surg Oncol* 2014;21:292–9.
- [50] Martin BM, Etra JW, Russell MC, et al. Oncologic outcomes of patients undergoing videoscopic inguinal lymphadenectomy for metastatic melanoma. *J Am Coll Surg* 2014;218:620–6.
- [51] Read RL, Haydu L, Saw RP, et al. In-transit melanoma metastases: incidence, prognosis, and the role of lymphadenectomy. *Ann Surg Oncol* 2015;22:475–81.
- [52] McMasters KM, Reintgen DS, Ross MI, et al. Sentinel lymph node biopsy for melanoma: controversy despite widespread agreement. *J Clin Oncol* 2001;19:2851–5.
- [53] Thalanayar PM, Agarwala SS, Tarhini AA. Melanoma adjuvant therapy. *Chin Clin Oncol* 2014;3:26.
- [54] Eggermont AM, Chiarion-Sileni V, Grob JJ, et al. Adjuvant ipilimumab versus placebo after complete resection of high-risk stage III melanoma (EORTC 18071): a randomised, double-blind, phase 3 trial. *Lancet Oncol* 2015.
- [55] Veenstra HJ, Wouters MW, Kroon BB, Olmos RA, Nieweg OE. Less false-negative sentinel node procedures in melanoma patients with experience and proper collaboration. *J Surg Oncol* 2011;104:454–7.
- [56] van der Ploeg IM, Kroon BB, Antonini N, Valdes Olmos RA, Nieweg OE. Is completion lymph node dissection needed in case of minimal melanoma metastasis in the sentinel node? *Ann Surg* 2009;249:1003–7.
- [57] Leiter U, Buettner PG, Eigentler TK, et al. Hazard rates for recurrent and secondary cutaneous melanoma: an analysis of 33,384 patients in the German Central Malignant Melanoma Registry. *J Am Acad Dermatol* 2012;66:37–45.
- [58] Pawlik TM, Ross MI, Thompson JF, Eggermont AM, Gershenwald JE. The risk of in-transit melanoma metastasis depends on tumor biology and not the surgical approach to regional lymph nodes. *J Clin Oncol* 2005;4588–90.
- [59] Veronesi U, Cascinelli N. Narrow excision (1-cm margin). A safe procedure for thin cutaneous melanoma. *Arch Surg* 1991;126:438–41.

- [60] Gadd MA, Coit DG. Recurrence patterns and outcome in 1019 patients undergoing axillary or inguinal lymphadenectomy for melanoma. *Arch Surg* 1992;127:1412–6.
- [61] Martini L, Brandani P, Chiarugi C, Reali UM. First recurrence analysis of 840 cutaneous melanomas: a proposal for a follow-up schedule. *Tumori* 1994;80:188–97.
- [62] Karakousis CP, Balch CM, Urist MM, Ross MM, Smith TJ, Bartolucci AA. Local recurrence in malignant melanoma: long-term results of the multiinstitutional randomized surgical trial. *Ann Surg Oncol* 1996;3:446–52.
- [63] Johnson RC, Fenn NJ, Horgan K, Mansel RE. Follow-up of patients with a thin melanoma. *Br J Surg* 1999;86:619–21.
- [64] Cohn-Cedermark G, Rutqvist LE, Andersson R, et al. Long term results of a randomized study by the Swedish Melanoma Study Group on 2-cm versus 5-cm resection margins for patients with cutaneous melanoma with a tumor thickness of 0.8–2.0 mm. *Cancer* 2000;89:1495–501.
- [65] Chao C, Wong SL, Ross MI, et al. Patterns of early recurrence after sentinel lymph node biopsy for melanoma. *Am J Surg* 2002;520–5.
- [66] Goydos JS, Patel KN, Shih WJ, et al. Patterns of recurrence in patients with melanoma and histologically negative but RT-PCR-positive sentinel lymph nodes. *J Am Coll Surg* 2003;196:196–204 [discussion 5].
- [67] Borgognoni L, Urso C, Vaggelli L, Brandani P, Gerlini G, Reali UM. Sentinel node biopsy procedures with an analysis of recurrence patterns and prognosis in melanoma patients: technical advantages using computer-assisted gamma probe with adjustable collimation. *Melanoma Res* 2004;14:311–9.
- [68] Macripo G, Quaglino P, Caliendo V, et al. Sentinel lymph node dissection in stage I/II melanoma patients: surgical management and clinical follow-up study. *Melanoma Res* 2004;14:S9–S12.
- [69] Thomas JM, Newton-Bishop J, A'Hern R, et al. Excision margins in high-risk malignant melanoma. *N Engl J Med* 2004;350:757–66.
- [70] Berk DR, Johnson DL, Uzieblo A, Kiernan M, Swetter SM. Sentinel lymph node biopsy for cutaneous melanoma: the Stanford experience, 1997–2004. *Arch Dermatol* 2005;141:1016–22.
- [71] Duprat JP, Silva DC, Coimbra FJ, et al. Sentinel lymph node biopsy in cutaneous melanoma: analysis of 240 consecutive cases. *Plast Reconstr Surg* 2005;115:1944–51 [discussion 52–3].
- [72] Nathansohn N, Schachter J, Gutman H. Patterns of recurrence in patients with melanoma after radical lymph node dissection. *Arch Surg* 2005;140:1172–7.
- [73] van Akkooi AC, de Wilt JH, Verhoef C, et al. Clinical relevance of melanoma micrometastases (0.1 mm) in sentinel nodes: are these nodes to be considered negative? *Ann Oncol* 2006;17:1578–85.
- [74] Cecchi R, De Gaudio C, Buralli L, Innocenti S. Lymphatic mapping and sentinel lymph node biopsy in the management of primary cutaneous melanoma: report of a single-centre experience. *Tumori* 2006;92:113–7.
- [75] Kretschmer L, Beckmann I, Thoms KM, Mitteldorf C, Bertsch HP, Neumann C. Factors predicting the risk of in-transit recurrence after sentinel lymphonodectomy in patients with cutaneous malignant melanoma. *Ann Surg Oncol* 2006;13:1105–12.
- [76] Dalal KM, Patel A, Brady MS, Jaques DP, Coit DG. Patterns of first-recurrence and post-recurrence survival in patients with primary cutaneous melanoma after sentinel lymph node biopsy. *Ann Surg Oncol* 2007;14:1934–42.
- [77] Roulin D, Matter M, Bady P, et al. Prognostic value of sentinel node biopsy in 327 prospective melanoma patients from a single institution. *Eur J Surg Oncol* 2008;34:673–9.
- [78] van den Broek FJ, Sloots PC, de Waard JW, Roumen RM. Sentinel lymph node biopsy for cutaneous melanoma: results of 10 years' experience in two regional training hospitals in the Netherlands. *In J Clin Oncol* 2013;18:428–34.
- [79] Ribero S, Osella-Abate S, Sanlorenzo M, et al. Favourable prognostic role of regression of primary melanoma in AJCC stage I-II patients. *Br J Dermatol* 2013;169:1240–5.
- [80] van der Ploeg AP, Haydu LE, Spillane AJ, et al. Outcome following sentinel node biopsy plus wide local excision versus wide local excision only for primary cutaneous melanoma: analysis of 5840 patients treated at a single institution. *Ann Surg* 2014;260:149–57.